

FLORIDA ENTOMOLOGIST

VOLUME 94

Instructions to authors

Your manuscript is scheduled to be printed in the next issue of **Florida Entomologist**. Attached are the preliminary proofs and a billing/Reprint Order form.

Please complete the following steps **within 48 hours**:

- 1) **Print out and Proof read the preliminary proofs.** Charges for author's alterations are \$1.25 per line; remember that a small change may often result in resetting several lines or an entire paragraph.

"Please note that the **Florida Entomologist** now includes **Key Words** for research reports and **Summaries** for scientific notes. In some cases, the Editor has added one or the other to the enclosed paper. You may make changes or additions you deem necessary on the preliminary proof."

- 2) **Examine the illustrations and captions.** Photocopies of line-drawings and half-tone screen prints will be the size and very near the quality that will appear in the published journal. **No original will be returned unless requested in writing to the Printer. Printer nor the Florida Entomologist Society will be responsible for plates and figures not reclaimed 30 days following publication.**

- 3) Describe your correction in an E-mail and send to:

Dr. Waldemar Klassen, Editor

editor.flagentsoc@gmail.com

- 4) All correspondence should be directed to Dr. Klassen not the printer.

Please Send Your Corrected Proofs and Manuscript to the Editor

SIGNS USED IN CORRECTING PROOFS

^	Insert marginal correction at this point in line	—	Less space
<i>tr</i>	Transpose; indicate by ~ in text	⊖	Close up entirely
<i>cap</i>	Capital; put 3 lines under letter or word	⊙	Period
<i>sc</i>	Small capital; put 2 lines under letter or word	^	Comma
<i>lc</i>	Lower-case letter; put oblique line through letter	⊙	Colon
<i>ital</i>	Italic; underline letter or word	;	Semicolon
<i>rom</i>	Roman letter; circle letter or word	ˆ	Apostrophe
<i>sp out</i>	Spell out; circle abbreviation	“ or ”	Quotation
<i>bf</i>	Boldface; underline letter or word with wavy line	=	Hyphen
<i>wf</i>	Wrong font; circle letter or word		Straighten lines
χ	Defective letter	[Move left
↓	Push down space]	Move right
⊗	Turn over	□	Em-quad space
⊖	Take out	$\frac{1}{m}$	One-em dash
^ v ^	Space evenly	¶	Make paragraph
#	Insert space	no ¶	No paragraph
□	Insert 2 spaces		

The signs listed above are used in marking proofs and they are readily understood by the printer. However, in the event of any possibility of ambiguity the instruction should be written fully on the proof rather than indicated by sign.

Courtesy of

E. O. Painter Printing Co., 4900 U.S. Highway 17 North, P.O. Box 877, DeLeon Springs, FL 32130 • Phone (386) 985-4877

Instructions for payment of Reprints & Society Charges

Florida Entomologist - Volume 94 - No. ____ (2011)

Return the billing forms and purchase order **IMMEDIATELY** to the **Printer:**

E. O. Painter Printing Company, P.O. Box 877, 4900 US Highway 17 North, DeLeon Springs, FL 32130

- *We cannot guarantee that reprint orders received more than 3 weeks after return of the preliminary proof will be filled.
- *Orders without attached purchase orders, prepayment or **completed** Visa or MasterCard form **will be refused**.
- *The charges for publication in **FLORIDA ENTOMOLOGIST** are: \$58.00 per page plus \$26.00 surcharge per figure or table.
- *There are no page charges for Book Reviews.
- ***Mailing Address** must include **street address**.

Instructions for payment

Purchase Orders - MUST INCLUDE the following or it will be returned.

- 1) A copy of a complete reprint order.
- 2) A copy of your sales tax exempt certificate (Florida P.O.).
- 3) Name, volume & number of journal **ON THE PURCHASE ORDER.**
- 4) Title of Article **ON THE PURCHASE ORDER.**
- 5) Author Names **ON THE PURCHASE ORDER.**
- 6) Should be made out to E. O. Painter Printing Company, Inc.

Credit Card Orders - MUST INCLUDE the following or it will be returned.

- 1) A copy of a complete reprint order and credit card order form.
- 2) A copy of your sales tax exempt certificate (Florida).
- 3) Name, volume & number of journal.
- 4) Title of Article.
- 5) Author Names.
- 6) Payment in U.S. funds from a U.S. Bank.

Check or Money Orders - MUST INCLUDE the following or it will be returned.

- 1) A copy of a complete reprint order.
- 2) A copy of your sales tax exempt certificate (Florida).
- 3) Name, volume & number of journal.
- 4) Title of Article.
- 5) Author Names.
- 6) Should be made out to E. O. Painter Printing Company, Inc.
- 7) Payment in U.S. funds from a U.S. Bank (including International Money Orders).

Florida Entomologist - Volume 94 - No. _____ (2011)

CAREFULLY COMPLETE THE BILLING/REPRINT ORDER FORM

Please check one:

_____ Purchase Orders _____ Check or Money orders (Complete this page and return this page by mail or fax to the printer.)
 _____ Credit Card Orders (Complete the following page and return this page by mail or fax back to the printer.)

Title: _____

Authors: _____

Ordering Reprints: _____ Yes or _____ No

Paying for Society Charges only: _____ Yes or _____ No

Reprint Price List

Minimum order is 100; reprints sold only in multiples of 100 per article. **No free reprints**

	Number of pages or fraction thereof								
Copies	1 - 4 pages	5 - 8 pages	9 - 12 pages	13 - 16 pages	17 - 20 pages	21 - 24 pages	25 - 28 pages	29 - 32 pages	33 - 36 pages
100	\$74.00	115.00	157.00	202.00	247.00	291.00	\$328.00	390.00	438.00
Add'l 100's	31.00	45.00	60.00	76.00	110.00	127.00	148.00	174.00	190.00

***Reprint orders mailed outside the contiguous U.S. (including Hawaii, Alaska, Canada, Mexico) Add \$40.00.**

_____ Copies		Research Papers	Scientific Notes	Worksheet: Use table above.
_____ No. of pages (Reprint charges from above chart)	\$ _____	\$ _____		
_____ Additional Postage \$40.00	\$ _____	\$ _____		
(For reprints shipped other than in the contiguous U.S.)				
_____ IFWA charge (enter \$100 or \$50) Include in total amount due:	\$100.00	\$50.00		
(The charge for Immediate Free Web Access is mandatory and is \$100 for articles and \$50 for scientific notes.)				
_____ No. of Society Page Charges @ \$58.00 each.	\$ _____	\$ _____		
_____ No. of Figures, Tables* @ \$26.00 each.	\$ _____	\$ _____		
*(Tables exceeding one page are charged per page)				
_____ No. of Authors Alterations @ \$ 1.25 each.	\$ _____	\$ _____		
_____ InfoLink	\$ _____	\$ _____		
(If you choose this option, add \$45.00 and fill out the attached "Request for InfoLink" form)				
_____ Florida State Sales Tax <i>Your county tax rate</i>	\$ _____	\$ _____		
_____ Total Amount Due:	\$ _____	\$ _____		

For purchases in the state of Florida, apply your county's local sales tax rate, including discretionary tax, to total amount due. Log on to www.myflorida.com/dor/taxes/ and click on *county tax rates* to find your county rate.

- All tax exempt purchases must supply a **copy of** a current Florida state sales tax exempt certificate DR 13 or 14 when submitting this form.
- All out of state purchases are exempt from sales tax.

E. O. Painter Printing Company, Inc.

* PO Box 877

* DeLeon Springs, FL 32130

* Fax: 386-985-4878

1. Mailing Address Information -Type or Print Clearly

(Physical address for U.S. deliveries)

First Name: _____ Last Name: _____

Business Name: _____

Address: _____

Phone: () _____ Fax: () _____ E-mail: _____

2. Billing Information -

First Name: _____ Last Name: _____

Business Name: _____

Address: _____

City: _____

State or Region: _____

Postal code: _____ Phone: () _____

Country: _____ Fax: () _____

COPY of the tax exempt certificate/ FL State Sales Tax Exempt Number (DR 13 or 14):

Credit Card Order Information

Complete & sign ALL four (4) Sections -- Fax to: 386-985-4878; Mail to: E. O. Painter Printing Company, P.O. Box 877, DeLeon Springs, FL 32130 or E-mail a signed scanned copy to: books@eopainterprinting.com

1. Mailing Address Information -Type or Print Clearly

(Physical address for U.S. deliveries)

First Name:

Last Name:

Business Name:

Address:

Phone: ()

Fax: ()

E-mail:

2. Billing Information - Your Billing Address is used to validate your Credit Card and should match the address on your Credit Card statement.

First Name:

Last Name:

Business Name:

Address:

City:

State or Region:

e-mail:

Postal code:

Phone: ()

Country:

Fax: ()

ALL tax exempt purchases **MUST send a **COPY** of the tax exempt certificate/ FL State Sales Tax Exempt Number (DR 13 or 14):**

3. Order Information-Type or Print Clearly

_____ **Copies** Florida Entomologist Volume 94 No.: _____, pages _____ to _____

Title _____

Senior Author: _____

Total Amount Due (Total Reprint charges from worksheet on reverse page) Charge to my credit card:\$_____

Name on the card _____

Select Credit Card: ☐ MasterCard ☐ VISA

				-					-					-			
--	--	--	--	---	--	--	--	--	---	--	--	--	--	---	--	--	--

				-		
--	--	--	--	---	--	--

Credit card number

3 Digits on the back of MasterCard/Visa

--	--	--

Expiration date

Signature _____ Date _____

4. Policy

*I agree that my credit card will be processed as a prepayment. Errors in my calculation of "total amount due" will result in this form being sent back for adjustment.

*I agree that reprints will be shipped four to six weeks after the publication has been mailed.

*I agree that delivery outside the contiguous United States can take four to six weeks.

*Mail, fax, or e-mail a **SIGNED** copy of this credit card order form.

Signature: _____

ESTABLISHMENT OF THE WEST INDIAN FRUIT FLY (DIPTERA: TEPHRITIDAE) PARASITOID *DORYCTOBRACON AREOLATUS* (HYMENOPTERA: BRACONIDAE) IN THE DOMINICAN REPUBLIC

COLMAR A. SERRA¹, MILEIDA FERREIRA¹, SOCORRO GARCÍA¹, LOENY SANTANA¹, MAIRA CASTILLO², CARIDAD NOLASCO³,
PAULA MORALES⁴, TIMOTHY HOLLER⁵, AMY RODA⁶, MARTIN ALUJA⁷ AND JOHN SIVINSKI⁸

¹Instituto Dominicano de Investigaciones Agropecuarias y Forestales (IDIAF) Santo Domingo,
Republica Dominicana

²Universidad Autónoma de Santo Domingo (UASD), Santo Domingo, Republica Dominicana

³Depto. de Sanidad Vegetal-Ministerio de Agricultura (DSV-MA), Santo Domingo, Republica Dominicana

⁴USDA-APHIS-IS, Ave. Pedro Henríquez Ureña, #133 - La Esperilla, Santo Domingo, Dominican Republic

⁵USDA-APHIS (ret.) at Center for Medical, Agricultural and Veterinary Entomology, 1600 SW 23rd Dr.,
Gainesville, FL 32608

⁶USDA-APHIS, Subtropical Horticultural Research Station, Miami, FL 33125

⁷Instituto de Ecología, A.C., Apartado Postal 63, 91000 Xalapa, Veracruz, México

⁸USDA-ARS, Center for Medical, Agricultural and Veterinary Entomology, 1600 SW 23rd Dr.,
Gainesville, FL 32608

ABSTRACT

The West Indian fruit fly, *Anastrepha obliqua* (Macquart), infests numerous fruit species, particularly Anacardiaceae and most importantly mango (*Mangifera indica* L.). Widespread in the Neotropics, it was first reported in Hispaniola nearly 70 years ago. Continental populations are attacked by the opiine braconid parasitoids *Utetes anastrephae* (Viereck) and *Doryctobracon areolatus* (Szépligeti). Largely sympatric, the two species co-exist through microhabitat specializations based on different ovipositor lengths and asymmetries in larval competitive abilities during multiparasitism. *Utetes anastrephae*, but not *D. areolatus*, is apparently native to the Dominican Republic. Since the two parasitoids share an evolutionary history over a substantial portion of their distributions it was proposed that 1) *D. areolatus* would find the Dominican environment suitable, as does *U. anastrephae*; and 2) that there would be no negative interactions when the two species were reunited and overall parasitism would increase. Immediately following releases, *D. areolatus* averaged 9% of adult insects recovered and two years after releases were concluded constituted a mean of 13%. By then the parasitoid had spread up to 50 km from release sites. There was no evidence of competitive exclusion of *U. anastrephae* by *D. areolatus*. Another opiine biological control candidate, *Diachasmimorpha longicaudata* (Ashmead), could be considered for release and establishment. Parasitoids alone are unlikely to provide economic levels of control, but can serve as components of an integrated pest management program established to maintain "fly-free" or "low prevalence" fruit export zones.

Key Words: *Anastrepha obliqua*, mango, biological control, invasive pest

RESUMEN

La mosca de las frutas de las Indias Occidentales, *Anastrepha obliqua* (Macquart), infesta numerosas especies de frutales, principalmente anacardiáceas, siendo el mango (*Mangifera indica* L.) especie de mayor importancia. Aunque este tefritido está ampliamente distribuido en el Neotrópico fue registrado hace 70 años en La Hispaniola. Las poblaciones continentales de *A. obliqua* son atacadas por los parasitoides braconídeos *Utetes anastrephae* (Viereck) y *Doryctobracon areolatus* (Szépligeti). Las dos especies de parasitoides co-existen a través de especializaciones basados en diferencias en la longitud del ovipositor y asimetrías en las habilidades competitivas durante el multiparasitismo. *Utetes anastrephae*, pero no así *D. areolatus*, es nativo en la República Dominicana donde ataca especies de *Anastrepha* nativas. Como las dos especies de parasitoides comparten una historia evolutiva sobre una porción sustancial de su rango de distribución, se pronosticó que: 1) *D. areolatus* se adaptaría al ambiente dominicano, tal como lo hace *U.*

anastrephae; y 2) no existirían interacciones negativas cuando las dos especies estén juntas y que el parasitismo total se incrementaría. Inmediatamente después de las liberaciones se recuperó un 9% de adultos de *D. areolatus* y dos años después de finalizadas las liberaciones se recobró un 13%. El parasitoide se diseminó por lo menos hasta 50 km de los sitios de liberación. No se observaron evidencias de exclusión competitiva de *U. anastrephae* por *D. areolatus*. Se recomienda a *Diachasmimorpha longicaudata* (Ashmead) como otro posible agente para el control biológico de *A. obliqua*. Los parasitoides por sí solos, probablemente no alcancen niveles económicos de control, pero pueden servir como componentes de un programa de manejo integrado de plagas para mantener zonas de exportación de frutas libres o de baja prevalencia de moscas.

Translation provided by the authors.

Neotropical and subtropical tephritid fruit flies infest hundreds of fruit and vegetable species (White & Elson-Harris 1992; Foote et al. 1993). Several polyphagous species are particularly destructive and responsible for trade restrictions wherever they occur (Siebert 1999; Siebert & Cooper 1995). *Anastrepha obliqua* (Macquart), the West Indian or Mango fruit fly, is one of the foremost of these pests and most frequently infests species of Anacardiaceae (Hernandez-Ortiz and Aluja 1993). *Spondias* spp. ("tropical plums") are particularly favored among native plants, but mango (*Mangifera indica* L.) is economically the most important host. Widespread, the fly occurs from the Greater (including Hispaniola) and Lesser Antilles to Mexico and south through Ecuador and parts of Brazil (Stone 1942; Foote et al. 1993). Wherever it exists, *A. obliqua* threatens both mangoes grown for local consumption, and complicates or precludes mango exports. Population levels in the Dominican Republic can be extremely high. For instance, up to 60 flies per trap have been captured in a single day in the mango orchards of Hato Damas (Serra et al. 2005a; Ogando & Serra 2006; Thomas et al. 2008). These high levels may be due to both a lack of sanitation measures and a depauperate natural enemy guild (Ovruski et al. 2000; Castillo et al. 2006).

The natural enemies of *A. obliqua* in Mexico have been investigated in detail (Sivinski et al. 1997, 1998; Aluja et al. 1998; Lopez et al. 1999), and 2 largely sympatric species of opiine braconids, *Utetes anastrephae* (Viereck) and *Doryctobracon areolatus* (Szépligeti), inflict high levels of mortality on *A. obliqua* in *Spondias* spp. Both *U. anastrephae* and *D. areolatus* attack late-instar *Anastrepha* spp. larvae, but because of a significantly shorter ovipositor, *U. anastrephae* is only efficient at reaching hosts in small fruit (Sivinski et al. 1998; 2001). *Doryctobracon areolatus* on the other hand is a major natural enemy of a variety of *Anastrepha* spp. larvae in a large number of fruit. In Florida, where biological control introductions have reunited the two species, only *D. areolatus* is found in *Psidium guajava* L. but both species occur in the much smaller *Eugenia uni-*

flora (Sivinski et al. 1998). Even within a fruit species the mean size of fruit containing *U. anastrephae* can be smaller than that of fruits containing *D. areolatus*. In spite of its shorter ovipositor, limited host range, lower fecundity and shorter adult life span (Sivinski et al. 2000), *U. anastrephae*, appears to persist because of its superior ability as an "intrinsic competitor" in instances of multiparasitism (Aluja et al., unpublished data).

Utetes anastrephae, but not *D. areolatus*, historically has parasitized native *Anastrepha* spp. on Hispaniola and occasionally parasitizes up to 90% of *A. obliqua* locally in *Spondias* spp. (Serra et al. 2005b). Because the two species co-occur commonly on the American mainland, it was predicted that 1) *D. areolatus* would find the Dominican environment suitable, as does *U. anastrephae*; and 2) there would be no negative interactions between 2 closely co-evolved species; and overall parasitism would increase with the introduction of *D. areolatus*, particularly among flies developing in larger fruits.

Two regions of the Dominican Republic were chosen for parasitoid releases: Hato Damas in the southern part of the country and Mata Larga in the northeast. Evidence of *D. areolatus* recovery, offspring of released parasitoids collected in the same fruiting season, and of establishment (*D. areolatus* collected in a subsequent fruiting season), are presented in the following. In addition, we discuss *U. anastrephae* parasitism, the effects of fruit size and host density on parasitism by both species and search for any indication of *D. areolatus* excluding *U. anastrephae* from hosts. The introduction of another opiine braconid species, *Diachasmimorpha longicaudata* (Ashmead), is considered and we argue that, in this particular case, sequential introductions might be more efficacious than simultaneous.

MATERIALS AND METHODS

Origin of Parasitoids

Doryctobracon areolatus were reared at the Instituto de Ecología, Xalapa, Veracruz, Mexico on

larvae of *Anastrepha ludens*. Details on the facilities, equipment and procedures are available in Aluja et al. (2009).

Release Sites

Parasitoids were periodically released at sites chosen on the basis of relatively high *Anastrepha* spp. densities and large numbers of *Spondias* spp., particularly *S. mombin* L. host trees in the Hato Damas area near San Cristóbal (18°24'59.51"N: 70°06'27.98"W) and in the Mata Larga area, and the rural surroundings of San Francisco de Macoris (19°18'01.83"N: 70°15'15.33"W). At each site there were initially 20 host tree-release points.

Release Schedule

Twenty one *D. areolatus* releases were made at approximately weekly intervals over the period, 17 Jun to 15 Dec 2005. Mortality and sex ratio of newly arrived parasitoids were recorded. They were then provided with honey and water, held for a min of 24 h, and separated into lots of 20 individuals (ca. 1 male: 2 females) each for transport to release sites. In total, 16,973 *D. areolatus*, 58% of them female, were taken into the field. Of these, 87% (14,791) were released in the Hato Damas region. Five releases totaling 2,182 (67% female) insects were made in the Mata Larga region from 22 Jul to 30 Sep 2005.

Fruit Sampling and Parasitoid Recovery

Freshly fallen fruit and, fruit from the canopy - when fallen fruits were absent - were collected, counted and weighed on the day of release, 2 days after and ~1 week after release. Sample sizes were based on estimates of available fruit (Sivinski et al. 1996; Sivinski et al. 1998). Larvae were allowed to emerge from fruit for a minimum of 10 d. They were then transferred to cups containing moistened sand as a pupation medium and held for another mo. Adult insect eclosion was noted over this period and insect identities were confirmed.

Statistical Analysis

With the most extensive data sets (insects from *S. mombin*; 2005, release year and 2007, a post-release year), the General Linear Models (GLM) procedure of the Statistical Analysis System (SAS Inst. 2004) was used to examine the following effects on parasitism by *U. anastrephae* and *D. areolatus*; estimated fruit size (= sample weight/number of fruit in the sample), mean numbers of hosts in the sample, mean parasitoid density (numbers of parasitoids / weight of fruit),

and parasitism by the other parasitoid species. Parasitism data, being proportional, was log-arc-sine transformed prior to analysis. Separate analyses were used to examine the effects of the following abiotic parameters on parasitism by both species: sample site (a class variable), Julian date and the quadratic of Julian date (to examine any possible non-linear relationships). In order to determine if parasitism by *D. areolatus* supplanted that of *U. anastrephae*, a Student t-test, with a Satterthwaite correction due to unequal variance, compared fruit samples that contained *U. anastrephae* and in which *D. areolatus* were either present or not. Lower mean parasitism by *U. anastrephae* in the presence of *D. areolatus* could indicate competition for hosts, but similar levels of *U. anastrephae* parasitism in the 2 categories would suggest that *D. areolatus* was providing additional parasitism.

RESULTS

Evidence of Establishment and Spread of *Doryctobracon areolatus*

In 2007, 2 years following parasitoid releases, *D. areolatus* was recovered from 17 of 18 of the monitored localities in the Hato Damas region. Where parasitism occurred, percent parasitism ranged from 1.1 to 100% (Fig. 1, derived from 18 *Spondias* spp. trees). In addition to being collected from *S. mombin*, *D. areolatus* was also collected from *Anastrepha* spp. in "tropical plum/ciruela" (*S. purpurea* L. var. 'flava' and var. 'purpurea', Anacardiaceae), golden apple/manzana de oro (*S. cytherea* Sonn., Anacardiaceae), guava/guayaba (*Psidium guajava* L., Myrtaceae), Malay apple/manzana malaya (*Syzygium malaccense* (L.) Merril & Perry, Myrtaceae), bilimbi/vinagrillo (*Averrhoa bilimbi* L., Oxalidaceae), carambola (*A. carambola* L., Oxalidaceae), and tropical almond/almendro tropical (*Terminalia catappa* L., Combretaceae). In the Mata Larga area establishment was finally confirmed by 2007 in collections of *S. mombin* (Table 1), although parasitism was typically less than 10%. This relatively low level may have been due to the lower number of released parasitoids and fewer alternative fruit fly hosts.

Following post-release collections at release sites in 2007, additional fruits were collected in 2007-2008 at distances of 2.5 to 50 km from the nearest point of release. *Doryctobracon areolatus* was obtained from *S. mombin*, *S. purpurea*, *A. carambola* and *P. guajava* (Table 2; Fig. 2). The farthest of these points was in a commercial guava (cv. 'Cubano') orchard located in Piedra Blanca, Monseñor Nouel province, 50 km north-east of the nearest release site in the Hato Damas area. We have no way of knowing if the parasitoid dispersed there on its own or was carried by humans. Additional data on emergence of fruit flies




Fig. 1. Pupal mortality and emergence of *Anastrepha* spp., *Doryctobracon areolatus* and *Utetes anastrephae* from fruits of 8 of the 16 local *Spondias mombin* trees and 2 *Spondius purpurea**, i.e., # 10 & 14, at Hato Damas, San Cristóbal, 2 Apr to 10 Oct 2007.

and parasitoids during a 3.5-mo period in 2008 are presented in Fig. 2. Besides one specimen of *U. anastrephae*, all recovered parasitoids were *D. areolatus*, which attacked up to 60% of the available hosts.

Effects of Biotic and Abiotic Variables and of Potential Competition on Parasitism

In 2005, during *D. areolatus* releases, 129 samples of *S. mombin* taken from 20 sites yielded 2958 fruits (mean = 22.9 [stderr = 1.4]) weighing a total of 26.8 kg (mean = 207.1 g [15.5]) and producing 16,074 *Anastrepha* sp. pupae, probably all *A. obliqua* (mean = 124.6 [11.0]). In 2007, a year following *D. areolatus* releases, 85 samples of *S. mombin* taken from 20 sites yielded 1625 pieces of fruit (mean = 19.1 [stderr = 1.1]) weighing a total of 16.3 kg (mean = 192 g [15.0]) and producing 4274 *Anastrepha* sp. pupae, probably all *A. obliqua* (mean = 50.3 [5.9]).

In 2005, mean parasitism (# parasitoid sp.₁ / # parasitoid sp.₁ + # parasitoid sp.₂ + # adult *A. obliqua*) by *U. anastrephae* was 0.34 (0.02), and 0.09 (0.01) by *D. areolatus*. In 2007, mean parasitism by *U. anastrephae* was 0.18 (0.02) and by *D. areolatus* 0.13 (0.02). Among sites, parasitism by *U. anastrephae* ranged from 0-1.0 in 2005 and from

0-0.90 in 2007. Among sites/dates, parasitism by *D. areolatus* ranged from 0-1.0 in 2005 and from 0-0.75 in 2007.

The following pertains only to insects obtained from *S. mombin* collected in 2005 (release year) and 2007 (2 years post-release). In 2005, during *D. areolatus* releases, parasitism by *U. anastrephae* was significantly and negatively related to mean estimated fruit size ($F = 5.5$, $df = 1$, $P < 0.02$). Parasitism by *D. areolatus* was not influenced by fruit size nor by host density or parasitism by *U. anastrephae* ($F = 1.63$, $df_{model} = 7$, $df_{error} = 121$, $P = 0.13$). It was subject to the effects of release sites ($F = 2.0$, $df = 12$, $P = 0.03$) and the interaction between release site and Julian date ($F = 2.1$, $df = 12$, $P = 0.03$). In 2007, subsequent to *D. areolatus* releases, there were no effects of fruit size, host density or parasitism by a competitor on either species ($F D. areolatus = 1.1$, $df_{model} = 7$, $df_{error} = 77$, $P = 0.40$; $F U. anastrephae = 1.0$, $df_{model} = 7$, $df_{error} = 77$, $p = 0.46$). Parasitism by *U. anastrephae* was affected by sample site ($F = 2.2$, $df = 1$, $P < 0.02$) and that by *D. areolatus* by Julian date ($F = 11.3$, $df = 1$, $P < 0.002$).

There was no evidence that *D. areolatus* excluded *U. anastrephae* from hosts during either year. In 2005 and 2007, there were no differences in *U. anastrephae* parasitism in samples with and

TABLE 2. COLLECTIONS OF FRUIT FROM NON-RELEASE SITES IN 2007-8 THAT YIELDED *DORYCTOBRACON AREOLATUS*. LOCATION REFERS TO THE DISTANCE (KM) AND DIRECTION FROM THE NEAREST RELEASE POINT.

Site, Host	Where fruit was collected Km and direction from nearest release site	Date	Number of fruit	Weight of fruits	Pupae/kg	# pupae	% Emergence			
							<i>Anastrepha</i>	<i>D. areolatus</i>	<i>U. Anastrephae</i>	Parasitism
La Cruz B, <i>S. mombin</i>	3,SW	21/08	25	155	97	15	68.0	4.0	28.0	32.0
La Cruz D, <i>S. mombin</i>	6,SW	1/10	25	198	250	40	100.0	0.0	0.0	0.0
Los Cocos, <i>S. mombin</i>	2.5,SE	3/09	85	335	314	85	45.9	2.4	51.8	54.1
Los Mameyes, <i>S. purpurea</i>	4.5,SE	21/08	16	400	100	40	90.0	10.0	0.0	10.0
Feliciano, <i>S. mombin</i>	5.5,SE	21/08	20	155	452	70	78.6	8.6	12.9	21.4
La Pared, <i>S. mombin</i>	7,SE	21/08	20	240	146	35	94.3	0.0	5.7	5.7
Piedra Blanca, <i>P. guajava</i>	50,NW	18/10	1	300	83	25	72.0	28.0	0.0	28.0

without *D. areolatus* ($t(2005) = 1.0$, $df = 66.8$, $P = 0.32$; $t(2007) = 1.4$, $df = 61$, $P = 0.15$).

DISCUSSION


Doryctobracon areolatus emerged from the puparia of *A. obliqua* following their release in the vicinity of fruiting *S. mombin*, *S. purpurea*, and *P. guajava* (infested with *A. suspensa*) and gave evidence of establishment when it was recovered from these same sites 2 years after releases had been concluded. Its ultimate distribution, density and capacity to control *A. obliqua* remain to be determined. This may not be the end of biological control efforts. Another candidate opiine parasitoid, *Diachasmimorpha longicaudata* (Ashmead), was not immediately released. It is an Indo-Malaysian exotic that has been introduced worldwide for tephritid control, and is a particularly effective natural enemy of *Anastrepha suspensa* (Loew) (Sivinski et al. 1996) We were concerned that it might suppress a critical alternative host for *D. areolatus* and so we postponed its establishment.

If *D. longicaudata* is also released the range of *D. areolatus* may shrink. In Mexico, *D. longicaudata* is more abundant than *D. areolatus* at higher altitudes (Sivinski et al. 2000) and in larger, commercial fruits (Lopez et al. 1999). It is unclear if the altitudinal distributions are the result of interspecific competition leading to habitat subdivision, but in Florida where first *D. areolatus* and then *D. longicaudata* was introduced to control *A. suspensa*, there is evidence of displacement (Eitam et al. 2004). Both species were originally established in southern Florida, but at present *D. areolatus* occurs in the northern portion of the peninsula (in spite of being more abundant at lower, warmer altitudes in Mexico) and *D. longicaudata* occurs in the southern portion with only a narrow range of overlap. One explanation for this distribution is that *D. areolatus* is better able to persist during periods of host scarcity and in areas with lower host-plant diversity. Thus the northern part of *A. suspensa*'s range provides a refuge from an otherwise superior competitor. In terms of biological control, parasitism levels were not noticeably affected by the replacement of one species by the other (Baranowski et al. 1993). There appears to be little reason for concern about the effects of *D. longicaudata* on the endemic *U. anastrephae*. It is widespread in Florida and occurs in the ranges of both introduced species (Eitam et al. 2004).

The possible effects of *D. areolatus*, a parasitoid with a broad host range within the genus *Anastrepha*, on native, non-target tephritids was also considered by the Ministry of Agriculture prior to release. In addition to *obliqua* and *suspensa* another 5 species of *Anastrepha* occur on

TABLE 1. RECOVERY OF *DORYCTOBRACON AREOLATUS* FROM *ANASTREPHA OBLIQUA* INFESTING *SPONDIAS MOMBIN* FRUITS COLLECTED IN THE MATA LARGA AREA IN 2007.

Tree #	Date	Pupae	% of emerged insects			
			<i>D. areolatus</i>	<i>U. anastrephae</i>	<i>Anastrepha</i>	Parasitism
2	21/8	70	1.4	11.4	87.1	12.9
3	21/8	39	2.6	15.4	82.1	17.9
4	21/8	65	3.1	0.0	96.9	3.1
4	3/9	32	9.4	6.3	84.4	15.6
9	3/9	69	6.3	25.0	68.8	31.3
9	26/9	32	5.0	0.0	95.0	5.0
10	26/9	24	6.0	0.0	94.0	6.0
12	21/8	44	4.5	4.5	90.9	9.1
15	21/8	83	7.3	0.0	92.7	7.3

Fig. 2. Infestation of *Psidium guajava* fruits, and the emergence of *Anastrepha* (probably *suspensa*) and braconid parasitoids at Piedra Blanca, Monseñor Nouel on 18 Oct 2007 (first detection) and between 25 Jan and 10 May 2008.

Hispaniola: *A. dissimilis* Stone, *A. insulae* Stone, *A. ocesia* Walker, *A. antillensis* Norrbom and *A. stonei* Steyskal (Stone 1942; Norrbom 1998). While there was no guarantee that *D. areolatus* would not attack these species as well, typically *D. areolatus* parasitism rates of uncommon, non-pestiferous flies are very low (<1%; e.g., Aluja et al. 2000; Aluja et al. 2003; Pereira et al. 2007; Rull et al. 2009).

Biological control, even with the addition of *D. areolatus*, is unlikely to unilaterally solve the economic problems created by *A. obliqua*. However, it could serve as a component of an integrated pest management program designed to protect "fly-free export" or "low-prevalence" zones, particu-

larly if high fly populations in non-commercial *Spondias* spp. could be suppressed leaving fewer flies to disperse into agricultural environments (Sivinski et al. 1996).

ACKNOWLEDGMENTS

We thank Dr. Robert Wharton, Department of Entomology, Texas A&M University who originally identified the parasitoids. Caroline Cohen USDA, APHIS-IS provided essential funding. Invaluable technical assistance was provided by Daniel Virgil, Eugenio Rodríguez, José Asiático, Pedro Marte, Raúl Nina, José Volquez (all DSV-MA), Xiomara Cayetano, Teofila Reinoso (IDIAF-CENTA), Lucía Silverio, Juan de Dios Moya, José Luis González y Luis Antonio Cabrera (ID-

IAF-EEML). Portions of this work were financially supported by INECOL, and by grants to MA from the Mexican Consejo Nacional de Ciencia y Tecnología (CONACyT Grant No. 46846-Q) and the Mexican Campaña Nacional contra Moscas de la Fruta (Secretaría de Agricultura, Ganadería, Desarrollo Rural y Pesca-Instituto Interamericano de Cooperación para la Agricultura (SAGARPA-IICA).

REFERENCES CITED

- ALUJA, M., HERRERA, E., LÓPEZ, M., AND SIVINSKI, J. 2000. First host plant and parasitoid record for *Anastrepha spatulata* Stone (Diptera: Tephritidae). *Proc. Entomol. Soc. Washington* 102: 1072-1073.
- ALUJA, M., LÓPEZ, M., AND SIVINSKI, J. 1998. Ecological evidence for diapause in four native and one exotic species of larval-pupal fruit fly (Diptera: Tephritidae) parasitoids in tropical environments. *Ann. Entomol. Soc. Am.* 91: 821-833.
- ALUJA, M., RULL, J., SIVINSKI, J., NORRBOM, A., WHARTON, R., MACIAS-ORDONEZ, R., DIAZ-FLEISCHER, F., AND LOPEZ, M. 2003. Fruit flies of the genus *Anastrepha* (Diptera: Tephritidae) and associated native parasitoids (Hymenoptera) in the Tropical Rainforest Biosphere Reserve of Montes Azules, Chiapas, Mexico. *Environ. Entomol.* 32: 1377-1385.
- ALUJA, M., SIVINSKI, J., OVRUSKI, S. M., GUILLEN, L., LOPEZ, M., CANCINO, J., TORRES-ANAYA, A., GALLEGOS-CHAN, G., AND RUIZ, L. 2009. Colonization and domestication of seven species of native New World hymenopterous larval-prepupal and pupal fruit fly (Diptera: Tephritidae) parasitoids. *Biocontrol Sci. Tech.* 19 (Suppl. No. 1): 49-79.
- BARANOWSKI, R., GLENN, H. AND SIVINSKI, J. 1993. Biological control of the Caribbean fruit-fly (Diptera, Tephritidae). *Florida Entomol.* 76: 245-251.
- CASTILLO, M., LÓPEZ, M., AND GUZMÁN, R. 2006. Prospección de moscas de las frutas en la República Dominicana. Reporte final para el Consejo Nacional de Investigaciones Agropecuarias y Forestales (CONIAF), Universidad Autónoma de Santo Domingo, (UASD), Santo. Domingo, República Dominicana.
- EITAM, A., SIVINSKI, J., HOLLER, T., AND ALUJA, M. 2004. Biogeography of the braconid parasitoids (Hymenoptera) of the Caribbean fruit fly, *Anastrepha suspensa* (Loew) (Diptera: Tephritidae). *Ann. Entomol. Soc. Am.* 97: 928-939.
- FOOTE, R., BLANC, F., AND NORRBOM, A. 1993. Handbook of the Fruit Flies (Diptera: Tephritidae) of America North of Mexico. Cornell University Press, Ithaca, New York.
- HERNÁNDEZ-ORTÍZ, V., AND ALUJA, M. 1993. Listado de especies del género Neotropical *Anastrepha* (Diptera: Tephritidae) con notas sobre su distribución y plantas hospederas. *Folia Entomol. Mexicana*: 88: 89-105.
- LOPEZ, M., ALUJA, M., AND SIVINSKI, J. 1999. Hymenopterous larval-pupal and pupal parasitoids of *Anastrepha* flies (Diptera: Tephritidae) in Mexico. *Biological Control* 15: 119-129.
- NORRBOM, A. L. 1998. A revision of the *Anastrepha daciiformis* species group (Diptera: Tephritidae). *Proc. Entomol. Soc. Washington*, 100: 160-192.
- OGANDO, F. R., AND SERRA, C. 2006. Desarrollo de un atrayente de material disponible en la República Dominicana para capturar Moscas de las Frutas. *Memoria Carib. Food Crop Soc. (CFCS)* 42: 102-112.
- OVRUSKI, S., ALUJA, M., SIVINSKI, J., AND WHARTON, R. 2000. Hymenopterous parasitoids on fruit-infesting Tephritidae (Diptera) in Latin America and the southern United States: Diversity, Distribution, Taxonomic Status and Their Use in Fruit Fly Biological Control. *Integrated Pest Manag. Rev.* 5:81-107.
- PEREIRA, R., STECK, G., VARONA, E., AND SIVINSKI, J. 2007. Biology and natural history of *Anastrepha interrupta* (Diptera: Tephritidae). *Florida Entomol.* 90: 389-391.
- RULL, J., WHARTON, R., FEDER, J. L., GUILLE, L., SIVINSKI, J., FORBES, A., AND ALUJA, M. 2009. Latitudinal variation in parasitoid guild composition and parasitism rates of North American hawthorn infesting *Rhagoletis*. *Environ. Entomol.* 38(3): 588-599.
- SAS, 2004. SAS/STAT User's Guide, version 9.1. SAS Institute, Cary, North Carolina.
- SERRA, C., GARCÍA, S., FERREIRA, M., BATISTA, O., EPSKY, N., AND HEATH, R. 2005a. Comparación de atrayentes para el trapeo de moscas de las frutas, *Anastrepha* spp. (Diptera: Tephritidae) en frutales en la República Dominicana. *Memoria Carib. Food Crop Soc. (CFCS)* 41(2): 524-532.
- SERRA, C., GARCÍA, S., AND FERREIRA, M. 2005b. Evaluación de *Mangifera indica*, *Spondias* spp. (Anacardiaceae) y *Psidium guajava* (Myrtaceae), hospederos de Moscas de las frutas, *Anastrepha* spp. (Diptera: Tephritidae), en cuanto a la presencia de parasitoides en diferentes zonas de la República Dominicana. Resúmenes. 51 Reunión Anual, Sociedad Interamericana para Horticultura Tropical (ISTH), 10-14/10/05, Boca Chica, República Dominicana.
- SIEBERT, J. 1999. Update on the economic impact of Mediterranean fruit fly on California agriculture. *Subtropical Fruit News* 7: 16-18.
- SIEBERT, J. B., AND COOPER, T. 1995. Embargo on California produce would cause revenue, job loss. *California Agriculture* 49: 7-12.
- SIVINSKI, J., ALUJA, M., HOLLER, T., AND EITAM, A. 1998. Phenological comparison of two braconid parasitoids of the Caribbean fruit fly (Diptera: Tephritidae). *Environ. Entomol.* 27: 360-365.
- SIVINSKI, J., ALUJA, M., AND LOPEZ, M. 1997. Spatial and temporal distributions of parasitoids of Mexican *Anastrepha* species (Diptera: Tephritidae) within the canopies of fruit trees. *Ann. Entomol. Soc. Am.* 90: 604-618.
- SIVINSKI, J., CALKINS, C., BARANOWSKI, R., HARRIS, D., BRAMBILA, J., DIAZ, J., BURNS, R., HOLLER, T., AND DODSON, G. 1996. Suppression of a Caribbean fruit fly (*Anastrepha suspensa* (Loew) Diptera: Tephritidae) population through augmented releases of the parasitoid *Diachasmimorpha longicaudata* (Ashmead) (Hymenoptera: Braconidae). *Biological Control* 6: 177-185.
- SIVINSKI, J., RENDÓN, P., HOLLER, T., BLOEM, K., MARTÍNEZ, A., AND LÓPEZ VALENZUELA, M. 2000. Distribution of hymenopterous parasitoids of *Anastrepha* spp. in guava (*Psidium guajava*) along an altitudinal transect in the Soconusco region (Chiapas, Mexico): practical implications for a potential augmentative biocontrol program with native parasitoids. 4th Mtg. Working Group on Fruit Flies of the Western Hemisphere, Mendoza, Argentina, Nov. 25-30. pp. 115-116.
- SIVINSKI, J., VULINEC, K., AND ALUJA, M. 2001. Ovipositor length in a guild of parasitoids (Hymenoptera: Braconidae) attacking *Anastrepha* spp. fruit flies (Diptera: Tephritidae) in Southern Mexico. *Ann. Entomol. Soc. Am.* 94: 886-895.

STONE, A. 1942. Fruit flies of the genus *Anastrepha*. United States Department of Agriculture Miscellaneous Publication, 72: 188-190.

THOMAS, D. B., EPSKY, N., SERRA, C., HALL, D., KENDRA, P., AND HEATH, R. 2008. Ammonia formulations

and capture of *Anastrepha* fruit flies (Diptera: Tephritidae). *J. Entomol. Sci.* 43(1): 76-85.

WHITE, I., AND ELSON-HARRIS, M. 1992. Fruit Flies of Economic Significance: Their Identification and Bionomics. C.A.B. International, Wallingford, UK.

PROOF